

## Sexually Transmitted Infections by Veronica Valles

Gonorrhea is a STI which is a sexually transmitted infection that is caused by Neisseria. The cause of this bacteria it can harm the mouth, and the rectum. For the women the uterus and the cervix is the first to which the infection will attack. This infection will go into the uterus harming many women when pregnant. This kind of infection can be found in teenagers of ages 15-19 and adults of 20-24 years of age.



The infection is spread the same as the HIV by intercourse. These two infections are very likely alike; having gonorrhea is the same as having HIV because of the gonorrhea causes the starting of the HIV. However, if pregnant gonorrhea if positive to a pregnant women then it could be passed to an infant while giving birth, but this will infect the baby's eyes the results of the baby getting gonorrhea is of sexual abuse.

Not being sure if having gonorrhea? Taking some tests that will look for some bacteria in the DNA. Checking for some discharged of a penis

and a cervix on a microscope slide with just a dye and looking for any bacteria. Most results are better for the men with quick results and not very helpful for the women. More often the doctors use urine or cervical swabs to do some tests. Some of other tests that are treated contain of use of discharge placed in a plate for 2 to 3 days for watching results of gonorrhea to grow. These samples could determine the infection of gonorrhea for 90 percent of the time. Other that has some test for looking gonorrhea in the throat has searching of drug use in bacteria.

For this kind of infection there are some treatments that are given to patients especially the most common treatment that are given are the one of prescribe of a hard and strong dose. Cefixime, Ceftriaxone, and Levofloxacin are some of the drugs use to calm or reduces the infection. There are some risks that some people take but more are the pregnant women and younger than 18 cannot take this kind of drug like Ciprofloxacin and Ofloxacin. Gonorrhea is just as common as chlamydial infection these two infections are around a person about the same time of each. In these cases the doctors prescribe an antibiotics combination for the two diseases.



There are also some difficulties when gonorrhea is not treated. This can cause the infection to become more than

it has started and infected more parts of your body to the point in which could kill you more quickly from the inside. While not controlling the gonorrhea it could go to the blood stream and infecting the joints and also the hearts valves.

Other treated infection is for the female that when treated its used of PID having some of a females tissue to form into a fallopian tubes this tube will fertilized the eggs inside and not them to pass the uterus to be occurred the results will be that the tube will be implant and causing a tubal pregnancy. If this becomes of some the results the complication is a death of a mother or a misarranged.



This infection can also be passed to an infant while being pregnant by the birth canal. Doctors can prevent the infection to your baby by immediately applying silver nitrate in the eyes of your new born after giving birth.